

Table XIV-10
LIVER AND INTRAHEPATIC BILE DUCT CANCER (Invasive)

SEER INCIDENCE AND U.S. MORTALITY
AGE-ADJUSTED RATES AND TRENDS^a
By Race/Ethnicity and Sex

SEER Incidence

RACE/ETHNICITY	Rate 2000-2004 ^b Rate per 100,000 persons			Trend 1995-2004 ^c APC (%)		
	Total	Males	Females	Total	Males	Females
	All Races	6.2	9.5	3.4	2.2*	2.4*
White	5.2	7.9	2.9	2.1*	2.3*	0.9
White Hispanic ^d	9.9	14.8	5.8	0.7	0.8	0.1
White Non-Hispanic ^d	4.6	7.1	2.6	1.9*	2.1*	0.4
Black	7.6	12.7	3.8	3.3*	4.5*	0.5
Asian/Pacific Islander	13.9	21.3	7.9	0.0	0.2	-0.1
Amer Ind/Alaska Nat ^e	9.7	14.8	5.5	0.2	-	-
Hispanic ^d	9.7	14.4	5.7	1.0	1.0	0.4

U.S. Mortality^f

RACE/ETHNICITY	Rate 2000-2004 Rate per 100,000 persons			Trend 1995-2004 APC (%)		
	Total	Males	Females	Total	Males	Females
	All Races	4.9	7.1	3.1	1.7*	1.8*
White	4.5	6.5	2.8	1.7*	1.9*	0.8*
White Hispanic ^d	7.9	11.2	5.2	1.4*	1.1*	1.3*
White Non-Hispanic ^d	4.2	6.1	2.7	1.5*	1.7*	0.5
Black	6.5	10.0	3.9	1.3*	1.7*	0.3
Asian/Pacific Islander	10.6	15.5	6.7	-0.6	-1.1*	0.4
Amer Ind/Alaska Nat						
Total U.S.	6.3	8.3	4.6	2.0	1.2	2.4
CHSDA Counties	8.4	10.7	6.4	2.2	1.5	2.7
Non-CHSDA Counties	3.5	5.1	2.2	-	-	-
Hispanic ^d	7.6	10.8	5.0	1.4*	1.1*	1.2

- The APC is the Annual Percent Change over the time interval.
- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.
- ^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^b Incidence data used in calculating the rates are from the 17 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).
- ^c Incidence data used in calculating the trends are from the 13 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia).
- ^d Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. The 2000-2004 Hispanic and Non-Hispanic death rates exclude deaths from Minnesota, New Hampshire and North Dakota. The 1995-2004 Hispanic and Non-Hispanic mortality trends exclude deaths from Maine, Minnesota, New Hampshire, North Dakota, and Oklahoma.
- ^e Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.
- ^f Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).
- * The APC is significantly different from zero (p<.05).